

Zetadex Gel Filtration Resin

For downstream purification of proteins and nucleic acids



Higher Resolution Narrower Band Widths Cleaner Separations

Zetadex is a beaded composite material developed by *emp BIOTECH* and comprised of ultrapure cross-linked dextran. It exhibits high selectivity, superb resolution, low non-specific adsorption and robust chemical stability.

Molecules purified with **Zetadex** are separated according to size. Smaller molecules pass significantly slower through the column than larger molecules.

The molecular weight cut-off (MWCO) for **Zetadex-25** is 5 kD for proteins and 10 bases for nucleic acids. For **Zetadex-50**, the cut-offs are 25 kD and 20 bases.

Buffer and pH effects on resolution are minimal and purified biomolecules are not significantly diluted (1.5-fold) when processed using **Zetadex**.

Zetadex is autoclavable at 121°C, pH 7 for 30 minutes and is stable in all commonly used buffers, including: 0.2M NaOH; 0.2M HCl; 1M acetic acid; 8M urea; 6M guanidine HCl; 1% SOS, 24% Ethanol; 30% Propanol; and 30% Acetonitrile.





ZetadexGel Filtration Resin

For downstream purification of proteins and nucleic acids

Gr	ade	Zetadex-25	Zetadex-50
Super Fine	Dry Bead Size:	20 – 50 μm (>80%)	20 – 50 μm (>80%)
	Water Regain:	2,15 – 2,25 ml/g	4.80 – 5.20 ml/g
	Swelling:	4 – 6 ml/g	9 – 11 ml/g
	MWCO (size exclusion):	below 5000 Da	below 25000 Da
	Fractionation Range:	1 – 5 kDa (globular proteins)	1 – 30 kDa (globular proteins)
	pH Stability:	2.0 to 13.0	2.0 to 13.0
	Catalog No.:	TM-0101	TM-0104
Fine	Dry Bead Size:	20 – 80 μm (>80%)	20 – 80 μm (>80%)
	Water Regain:	2,15 – 2,25 ml/g	4.80 – 5.20 ml/g
	Swelling:	4 – 6 ml/g	9 – 11 ml/g
	MWCO (size exclusion):	below 5000 Da	below 25000 Da
	Fractionation Range:	1 – 5 kDa (globular proteins)	1 – 30 kDa (globular proteins)
	pH Stability:	2.0 to 13.0	2.0 to 13.0
	Catalog No.:	TM-0102	TM-0105
Medium	Dry Bead Size:	50 – 150 μm (>80%)	50 – 150 μm (>80%)
	Water Regain:	2,15 – 2,25 ml/g	4.80 – 5.20 ml/g
	Swelling:	4 – 6 ml/g	9 – 11 ml/g
	MWCO (size exclusion):	below 5000 Da	below 25000 Da
	Fractionation Range:	1 – 5 kDa (globular proteins)	1 – 30 kDa (globular proteins)
	pH Stability:	2.0 to 13.0	2.0 to 13.0
	Catalog No.:	TM-0103	TM-0106
Coarse	Dry Bead Size:	100 – 300 μm (>80%)	100 – 300 μm (>80%)
	Water Regain:	2,15 – 2,25 ml/g	4.80 – 5.20 ml/g
	Swelling:	4 – 6 ml/g	9 – 11 ml/g
	MWCO (size exclusion):	below 5000 Da	below 25000 Da
	Fractionation Range:	1 – 5 kDa (globular proteins)	1 - 30 kDa (globular proteins)
	pH Stability:	2.0 to 13.0	2.0 to 13.0
	Catalog No.:	TM-0112	TM-0113
Agglutination Grade (Gel Card)	Dry Bead Size:	Not applicable 20 – 50 μm (>93%) 4.80 – 5.20 ml/g 9 – 11 ml/g TM-0111	20 – 50 μm (>93%)
	Water Regain:		4.80 – 5.20 ml/g
	Swelling:		
	Catalog No.:		
Pre-hydrated and Ready-to-use		Zetadex-25 Medium	Zetadex-50 Fine
	Hydration Buffer:	Hydrated in phosphate buffered saline; pH 7.4, w	ith 0.2% NaN ³ Hydrated in phosphate buffered saline; pH 7.4, with 0.2% N
	Catalog No.:	TM-0107	TM-0108
	Hydration Buffer:	Hydrated in deionized water with 0.15% ProCline	Hydrated in deionized water with 0.15% ProCline
	Catalog No.:	TM-0114	TM-0115







CHROMATOGRAPHY

ZetadexAntibody Purification

Using Zetadex-25 Medium Gel Filtration Resin

Sample: Mouse IgG in PBS

Column: Zetadex-25 M in BPG

300/500

Conc.: 12.5 mg/ml

Diameter: 30 cm

Volume: 4500 ml

Gel Volume: 25 Liters

Total IgG: 56.25 grams

Pressure: 1.0 bar (0.1 mPa or 13 psi)

Buffer: 100 mM Glycine-HCl, pH 2.0 neutralised with 1 M

Tris-HCl pH 8.0

System: ÄktaPilot (GE Healthcare)

Flow rate during


sample loading: 400 ml

Flow rate during equilibration

and elution: 500 ml

Total run time: 63 min

High Performance Results:



For more information, please contact us at +49 30 9489 2201 or tech@empbiotech.com

